arXiv:1511.04570v1 [math.CV] 14 Nov 2015

A SHORT PROOF OF CARTAN'S NULLSTELLENSATZ FOR ENTIRE FUNCTIONS IN $\mathbb{C}^n$

RAYMOND MORTINI

Abstract. Using the fact that the maximal ideals in the polydisk algebra are given by the kernels of point evaluations, we derive a simple formula that gives a solution to the Bézout equation in the space of all entire functions of several complex variables. Thus a short and easy analytic proof of Cartan's Nullstellensatz is obtained.

1. Introduction

The aim of this note is to give a short and easy proof of Cartan's Nullstellensatz:

Theorem 1.1. *Let $H(\mathbb{C}^n)$ be the space of functions holomorphic in $\mathbb{C}^n$. Given $f_j \in H(\mathbb{C}^n)$, the Bézout equation $\sum_{j=1}^N g_j f_j = 1$ admits a solution $(g_1, \dots, g_N) \in H(\mathbb{C}^n)^N$ if and only if the functions f_j have no common zero in $\mathbb{C}^n$.*

The usual proofs use a lot of machinery from sheaf theory, cohomology, (see for example [2]), or are based on the Hörmander-Wolff method by solving higher order $\overline{\partial}$-equations using the Koszul complex, a tool from homological algebra (see [6, p. 128-131]). The one-dimensional case, first done by Wedderburn, is very easy (see for example [4, p. 118-120] for the classical approach or [1, p. 130] for the $\overline{\partial}$-approach). For our proof to work, we shall only use a standard fact from an introductory course to functional analysis, namely Gelfand's main theorem: the maximal ideals in a commutative unital complex Banach algebra coincide with the kernels of the multiplicative linear functionals (see for instance [5]). The idea is to apply this result to the polydisk algebras on an increasing sequence of polydisks $\mathbf{D}_k$ and to glue together the solutions to the Bézout equations $\sum_{j=1}^N g_j f_j = 1$ on $\mathbf{D}_k$ by using a Mittag-Leffler type trick. The major hurdle to overcome was of course to find suitable summands that guarantee at the end the holomorphy.

Date: June 1, 2015.

1991 *Mathematics Subject Classification.* Primary 32A15; Secondary 46J15.

Key words and phrases. Entire functions; Cartan's Nullstellensatz; polydisk algebra; maximal ideals.

2. The general solution to the Bézout equation

Let $\mathbb{D}$ be the unit disk and let $A(\mathbf{D}^n)$ be the polydisk algebra; that is the algebra of those continuous functions on the closed polydisk

$$\mathbf{D}^n = \{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_j| \leq 1\}$$

which are holomorphic in $\mathbb{D}^n$. Endowed with the supremum norm, $A(\mathbf{D}^n)$ becomes a uniform algebra and coincides with the closure on $\mathbf{D}^n$ of the polynomials in $\mathbb{C}[z_1, \dots, z_n]$. We actually only need that $A(\mathbf{D}^n)$ is the uniform algebra generated by the coordinate functions Z_j, $j = 1, \dots, n$ on $\mathbf{D}^n$. It is now straightforward to show that an ideal I in $A(\mathbf{D}^n)$ is maximal if and only if it coincides with $M(a_0) = \{f \in A(\mathbf{D}^n) : f(a_0) = 0\}$ for some $a_0 \in \mathbf{D}^n$ (just take a character m on $A(\mathbf{D}^n)$ and put $a_0 = (m(Z_1), \dots, m(Z_n))$). Hence the Bézout equation $\sum_{j=1}^N x_j f_j$ in $A(\mathbf{D}^n)$ has a solution if and only if $\bigcap_{j=1}^N Z_{\mathbf{D}^n}(f_j) = \emptyset$, where $Z_{\mathbf{D}^n}(f)$ is the zero set of f on $\mathbf{D}^n$. We will use the following well-known elementary result. For the reader's convenience we reproduce the proof here (see [3]), because its understanding is fundamental for our construction.

Lemma 2.1. *Let R be a commutative unital ring. Suppose that $\mathbf{a} = (a_1, \dots, a_N)$ is an invertible N-tuple in R^N and let $\mathbf{x} = (x_1, \dots, x_N)$ satisfy $\sum_{j=1}^N x_j a_j = \mathbf{1}$; that is $\mathbf{x}\,\mathbf{a}^t = \mathbf{1}$. Then every other representation $\mathbf{1} = \sum_{j=1}^N y_j a_j$ of $\mathbf{1}$ can be deduced from the former by letting $\mathbf{y} = \mathbf{x} + \mathbf{a}H$, where H is an antisymmetric $(N \times N)-$matrix over R; that is $H = -H^t$, where H^t is the transpose of H.*

Proof. Suppose that $\mathbf{1} = \mathbf{x}\mathbf{a}^t$ and $\mathbf{1} = \mathbf{y}\mathbf{a}^t$. For $k = 1, \dots, N$, multiply the first equation by y_k and the second by x_k. Then

$$x_k - y_k = \sum_{j \neq k} a_j (y_j x_k - y_k x_j).$$

Thus $\mathbf{y} = \mathbf{x} + \mathbf{a}H$ for some antisymmetric matrix H.

To prove the converse, let $\mathbf{1} = \mathbf{x}\mathbf{a}^t$. Since H is antisymmetric we have (due to the transitivity of matrix multiplication and $\mathbf{x}\mathbf{y}^t = \mathbf{y}\mathbf{x}^t$)

$$\begin{aligned} (\mathbf{a}H)\mathbf{a}^t &= \mathbf{a}(H\mathbf{a}^t) = \mathbf{a}(\mathbf{a}H^t)^t \\ &= \mathbf{a}(-\mathbf{a}H)^t = (-\mathbf{a}H)\mathbf{a}^t. \end{aligned}$$

Thus $(\mathbf{a}H)\mathbf{a}^t = 0$. Hence

$$\mathbf{y}\mathbf{a}^t = (\mathbf{x} + \mathbf{a}H)\mathbf{a}^t = \mathbf{x}\mathbf{a}^t + (\mathbf{a}H)\mathbf{a}^t = \mathbf{1} + 0 = \mathbf{1}.$$

□

3. Proof of Theorem 1.1

Proof. Let $f_j \in H(\mathbb{C}^n)$ and put $\mathbf{f} = (f_1, \dots, f_N)$. Suppose that $\bigcap_{j=1}^N Z(f_j) = \emptyset$. For $k \in \mathbb{N}^*$, let $\mathbf{D}_k = (k\,\mathbf{D})^n$ be the closed polydisk

$$\mathbf{D}_k = \{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_j| \leq k\}.$$

Note that $\mathbf{D}_k \subseteq \mathbf{D}_{k+1}^{\circ}$. Let $\mathbf{a}_k \in A(\mathbf{D}_{k+1})^N$ be a solution to the Bézout equation $\mathbf{a}_k \cdot \mathbf{f}^t = 1$ on $\mathbf{D}_{k+1}$. Using Tietze's extension theorem [1], we may assume that the tuples $\mathbf{a}_k$ have been continuously extended to $\mathbb{C}^n$ [2]. By Lemma 2.1, there is an antisymmetric matrix H_k over $A(\mathbf{D}_{k+1})$ such that

$$\mathbf{a}_{k+1} = \mathbf{a}_k + \mathbf{f} \cdot H_k.$$

Put $\mathbf{a}_0 = \mathbf{0}$ and $H_0 = \mathbf{O}$. For $k = 0, 1, \ldots$, let P_k be an antisymmetric $N \times N$-matrix of polynomials in $\mathbb{C}[z_1, \ldots, z_n]$ such that

$$\max_{\mathbf{D}_{k+1}} ||\mathbf{f} \cdot H_k - \mathbf{f} \cdot P_k||_N < 2^{-k}.$$

We claim that the N-tuple

$$\mathbf{g} := \sum_{k=0}^{\infty} \big(\mathbf{a}_{k+1} - \mathbf{a}_k - \mathbf{f} \cdot P_k\big)$$

belongs to $H(\mathbb{C}^n)^N$ and is a solution to the Bézout equation $\mathbf{g} \cdot \mathbf{f}^t = 1$ in $H(\mathbb{C}^n)$. In fact, let $\mathbf{D}_m$ be fixed. Then the series defining $\mathbf{g}$ is uniformly convergent on $\mathbf{D}_m$ since

$$\begin{aligned} \mathbf{g} &= \sum_{k=0}^{m} \big(\mathbf{a}_{k+1} - \mathbf{a}_k - \mathbf{f} \cdot P_k\big) + \sum_{k=m+1}^{\infty} \big(\mathbf{a}_{k+1} - \mathbf{a}_k - \mathbf{f} \cdot P_k\big) \\ &= \mathbf{a}_{m+1} - \mathbf{f} \cdot \big(\sum_{k=0}^{m} P_k\big) + \sum_{k=m+1}^{\infty} \big(\mathbf{a}_{k+1} - \mathbf{a}_k - \mathbf{f} \cdot P_k\big) \end{aligned}$$

and the tail can be majorated on $\mathbf{D}_m$ by

$$\sum_{k=m+1}^{\infty} \|\mathbf{f} \cdot H_k - \mathbf{f} \cdot P_k||_N < 2^{-m}.$$

Moreover, on $\mathbf{D}_m$, $\mathbf{a}_{m+1}$ and all the summands in the series $\sum_{k=m+1}^{\infty}$ are holomorphic. Since m was arbitrarily chosen, we conclude that $\mathbf{g} \in H(\mathbb{C}^n)^N$. Moving again to $\mathbf{D}_m$ we see that, due to the antisymmetry of the matrices H_k and P_k,

$$\begin{aligned} \mathbf{g} \cdot \mathbf{f}^t &= \mathbf{a}_{m+1} \cdot \mathbf{f}^t - \mathbf{f} \cdot \big(\sum_{k=0}^{m} P_k\big) \cdot \mathbf{f}^t + \sum_{k=m+1}^{\infty} \mathbf{f} \cdot (H_k - P_k) \cdot \mathbf{f}^t \\ &= 1 - 0 + 0 = 1. \end{aligned}$$

□

[1] Since we will consider a telescoping series $\sum T_j$, where the domains of definition of the summands T_j are strictly increasing, even an application of Tietze's theorem is not necessary.

[2] Of course, outside $\mathbf{D}_k$ the Bézout equation is not necessarily satisfied.

Acknowledgements

I thank the referee of the journal "American Math. Monthly" for some useful comments concerning the introductory section of a previous version of the paper.

References

[1] M. Andersson. *Topics in Complex Analysis*, Springer, New York 1997. 1

[2] S. Krantz, *Function theory of several complex variables*, Reprint of the 1992 edition. AMS Chelsea Publishing, Providence, RI, 2001. 1

[3] R. Mortini, B. Wick, Simultaneous stabilization in $A_{\mathbb{R}}(\mathbb{D})$, Studia Math. 191 (2009), 223–235. 2

[4] R. Remmert. *Funktionentheorie II*, Springer, Berlin , 1991. 1

[5] W. Rudin, *Real and Complex Analysis*, third edition, McGraw-Hill, New York, 1986 1

[6] E. Sawyer, *Function Theory: Interpolation and Corona Problems*, Fields Institute Monographs, Amer. Math. Soc. 2009. 1

Université de Lorraine, Département de Mathématiques et Institut Élie Cartan de Lorraine, UMR 7502, Ile du Saulcy, F-57045 Metz, France

E-mail address: raymond.mortini@univ-lorraine.fr